
wartpeel®
(Patented Compounded 5Fu, sal acid)

WartPEEL® Instructions for Use FOR SEBORRHEIC KERATOSIS

PRIOR TO USING THIS MEDICATION, IT IS IMPORTANT TO INFORM YOUR PRESCRIBER IF YOU ARE BREASTFEEDING, PREGNANT, OR THINKING OF BECOMING PREGNANT. DO NOT USE IF PREGNANT, TRYING TO GET PREGNANT OR BREASTFEEDING.

STEP 1: Apply once daily at bedtime. Remove tip cap and apply a small amount of WartPEEL directly onto the plastic applicator (or a toothpick). Use forward and backward pressure on the plunger to control the amount of medication that comes out of the tip of the syringe. Replace tip cap to prevent the medication from drying out.



STEP 2: WartPEEL is a glue not a cream, so make sure it fully dries. After applying a thin and even film on the wart or other lesions being treated, wait until it dries completely (10 to 20 minutes) and a color change from white to translucent white occurs. Test dryness with the blue applicator. WartPEEL should not stick to the applicator and is slightly soft when fully dry. The drying process can be sped up by using a hair dryer on the cool setting or a fan. Be careful not to get the medication onto the healthy skin as this may lead to irritation.

STEP 3: Tape may come with your WartPEEL prescription but is not recommended when treating seborrheic keratosis.

STEP 4: Wash hands after applying the medication. Never apply to the mouth, eyes, or inside the nose.

STEP 5: In the morning wash the area to remove excess medication. The area can be left uncovered during the day or a covering such as a band aid may be used.

STEP 6: Repeat steps 1-5, applying and removing the medication once daily for 5-10 days or as directed.

Make sure to view our video tutorial for more detailed instructions, tips and real life case studies to help make your WartPEEL treatment is successful!
CLICK HERE - SEBORRHEIC KERATOSIS FOR PATIENTS

Please see following information on frequently asked questions and storage information.

www.wartPEEL.com

wartpeel®
(Patented Compounded 5Fu, sal acid)

Frequently Asked Questions (FAQs)

Will my prescription insurance pay for WartPEEL?
Since WartPEEL is a compounded medication that is custom made, most insurance companies do not cover it. In many cases, the prescription is eligible to be paid for using a Flex Spending Account or a Health Savings Account. The patient may also choose to self-submit a claim. If you would like to do this, please contact your insurance company for further instructions.

What will I receive in the package from the compounding pharmacy?
You will receive a receipt and a prescription vial. The vial contains a syringe with the medication, two blue applicators, and a roll of Blenderm tape (if requested by your provider).

What should I expect when starting WartPEEL?
After the first few days of application, the wart will usually turn white in color if it is in a thick-skinned area like the hands, feet, elbows, and knees. In thin skinned areas like the arms, legs, trunk, face and genitals, the warts will at first swell and then form scabs. Please see our patient instruction sheets with video links for more information on treating different types of skin lesions on wartpeel.com.

Should I continue using the medication until the syringe is empty?
The prescribed quantity is a standard amount that may not be completely used over the course of treatment.

How do I know when to stop treating?
The average treatment time for most patients is 1 to 3 weeks. Appearance of the lesion being treated over the course of therapy varies. See the case studies at the bottom of the instructions for the type of lesions you are treating at wartpeel.com. There are several real-life examples of treating various types of skin lesions. A follow-up appointment with a prescriber may be necessary to determine when to discontinue.

What are the most common side effects?
Redness and irritation around the wart(s) are common and can be minimized with careful application to the wart, avoiding the surrounding skin. Mild to moderate burning and soreness can be expected especially towards the end of the treatment. This means WartPEEL is penetrating from the epidermis; the top layer of the skin where the wart lives, to the dermis; the layer underneath that is bright red and has many nerves. Though uncommon, additional adverse effects include allergic contact dermatitis and itchiness. There are no known incidences of scarring to date. It is believed the WartPEEL is not strong enough to scar the skin. Call your doctor or get medical help if you have any side effects that bother you or do not go away.

Can I leave the medication and tape on for 24 hours or do I have to take it off in the morning?
Remove the tape in the morning and wash off any visible WartPEEL.

Do I have to use the tape provided or can I use another type of tape such as duct tape?
We supply a roll of Blenderm tape made by 3M because it is latex free and enhances the effectiveness of WartPEEL. Duct tape may be used if desired based on individual preference or prescriber preference. Tape should only be used on the hands, feet, elbows and knees.

Can I apply the medication directly to the wart or must I use the blue applicator?
You should use the blue applicators, or another utensil such as a Q-tip or toothpick, to prevent any direct contact between the wart and syringe tip because it can contaminate the medication in the syringe.

How do I clean the blue applicators?
You should wash them with soap and water after each use.

How do I store the medication?
Store this medication in the prescription vial at a controlled room temperature out of direct sunlight. It should be stored out of the reach of children.

How long is the medication good for?
WartPEEL should not be used past the beyond-use date. Doing so will not give the desired outcome and may result in potentially harmful effects.

How do I dispose of this medication?
Refer to your individual state's regulations for drug disposal. Many local law enforcement agencies participate in "take-back" days several times per year. These local events and other local permanent collection programs can be found on the DEA website.

Is WartPEEL FDA approved?
The individual active components of WartPEEL (salicylic acid and 5-fluorouracil) are FDA approved and have been used to treat warts and other skin conditions for decades. The combination product is a compounded medication developed exclusively by MedCara and produced under the strictest compounding regulations. Compounded medications like WartPEEL do not require and are not given FDA approval. For further information, watch a pharmacist discuss the product in more detail in our Patient Education Video at wartpeel.com.

Why should WartPEEL be kept out of the reach of pregnant or nursing women?
WartPEEL has not been studied in pregnant or nursing women so it is not recommended these populations be exposed to the medication. Although previous studies have shown minimal medication will be absorbed into the body when applied on the wart, the product contains 5-fluorouracil which can cause birth defects when used in higher doses. It is important to keep this product out of direct contact with pregnant or nursing mothers. If there is not another individual available to apply the medication to the patient, pregnant/nursing individuals should wear gloves and wash their hands thoroughly after assisting with application.

What is the minimum age that a child must be to use WartPEEL?
The ingredients in WartPEEL have been studied in patients as young as 4 years old, but some health care providers have used the medication in younger patients. Speak with your dermatologist if your child is under 4 years old.

Who do I contact if I have additional questions about WartPEEL?
For questions on WartPEEL, feel free to use the toll-free number listed on your prescription label to contact us over the course of your treatment. Additional resources and training videos are available on Wartpeel.com.

What makes WartPEEL different?
WartPEEL is a highly effective prescription medication that has three US patents and has been prescribed and recommended by dermatologists, podiatrists, and pediatricians since 2004. Most topical wart treatments fail because they do not penetrate through the thick layers of dead skin that cover the wart. They also spread to the normal skin around the wart which causes severe irritation. WartPEEL is not a cream or ointment, but an adhesive gel specifically designed to penetrate the many layers of dead skin to get down to the level of the living wart. See what other health care providers are saying about WartPEEL at wartpeel.com.

How much does WartPEEL cost?
$98 with shipping included (5 to 7 days).
Do I rub WartPEEL into the skin?
No! WartPEEL is an adhesive gel similar to a glue. You should apply it as precisely as possible to the wart and let it fully dry. This usually takes around 10 to 15 minutes. Drying time can be accelerated with a fan. When fully dry, it is slightly soft and dry to the touch.

Why can't WartPEEL be sent to my local pharmacy?
WartPeel is a patented compounded medication licensed only to be produced by NuCara Pharmacy where it has been made since 2004.

How long will it take to receive WartPEEL?
Once we receive payment, it will take around 5 to 7 days to receive the medication. The pharmacy will attempt to contact you within 48 hours via phone call and/or text message.

I looked up my pharmacy and it's not in my state. Is that a mistake?
No. NuCara pharmacies have licenses to ship to 48 states but are only located in three states. The pharmacy that ships to your state will most likely be in another state.

The pharmacy hasn't called me, what do I do?
The vast majority of the time, the pharmacy does contact you within 24 hours by phone or text but isn't always successful. Don't hesitate to give the pharmacy a call to expedite the process.

How do I get refills of WartPEEL?
Your prescriber may have added refills on the prescription. You can locate this information on the vial that contains the medication. If you do have refills, you can contact the compounding pharmacy using the phone number listed on the label to ask for a refill. If your prescriber did not include refills, our pharmacy can contact them to
request refills, or you can contact your health care provider to ask for a refill.

When do I apply the medication?
WartPEEL is applied at night and washed off in the morning. Skipping days is not recommended as it lengthens treatment times and may cause treatment failure.

What do I do with the buildup of all the white stuff?
White stuff is good! It is part of a dead wart and is recommended to be removed in a safe and sterile way. This can be done with a nail file, cuticle pusher, pumice stone, tweezers or the WartPEEL applicator. Do not use anything sharp and stop if you experience pain. The more the white debris is removed, the closer the WartPEEL is to reaching the wart and the better it works. Sterilize tools between use and discard any tool that cannot be sterilized after the treatment is over to avoid spreading warts to other areas.

Storage Requirements
Store this medication at room temperature. Keep out of reach of children.

Expiration Date
The medication is best used within four months of the date it was made. Do not use or keep outdated medication.

Side Effects
- Redness and irritation around the wart(s) are most commonly reported
  - Irritation can be minimized with careful application to the wart alone
  - Irritation can be minimized by allowing the medication to dry completely before applying tape • Though uncommon, burning, crusting, allergic contact dermatitis, itchiness, scarring, rash, soreness, and ulceration can occur.
- Call your doctor or seek medical attention if you have any side effects that bother you or do not go away.

Contraindications
- Do not use if you are pregnant or breastfeeding
  - May cause fetal harm when given to pregnant women
  - Medications present in breast milk may lead to life-threatening or fatal side effects in nursing babies
- Do not use if you have a severe allergic reaction (anaphylaxis) to WartPEEL or its components including:
  - Salicylic acid or derivatives such as aspirin
  - 5-fluorouracil (5-FU)